

Physics Course - Exercises Summer 2009

Stefan Kremer (kremer@tkm.uni-karlsruhe.de)

Extra Tutorial

Holger Schmidt (hschmidt@tkm.uni-karlsruhe.de)

3.10.2009

1. Exercise

Consider two masses $m_1 > m_2$ on opposite sites of a roof with angle α with respect to the ground. Both masses are at height h and are connected by a rope over the top of the roof. Both masses are exposed to gravitational force ($g = 10 \frac{\text{m}}{\text{s}^2}$).

- (i) Determine the acceleration of the system.
- (ii) Find the time t_1 the mass m_1 hits the ground.
- (iii) What will be the tension force in the rope?
- (iv) Calculate these quantities for $\alpha = 30^\circ$, $m_1 = 8 \text{ kg}$, $m_2 = 4 \text{ kg}$ and $h = 10 \text{ m}$.

2. Exercise

A uniform rod of mass m and length L is hung at a distance $\frac{L}{4}$ of one of its ends and is initially pulled back at an angle $\phi = \phi_0$ to the vertical. The rod is exposed to gravity. Vertical below the attachment a spring with constant D is placed.

- (i) What is the moment of inertia of the system?
- (ii) Determine the angular velocity when the rod reaches the vertical position.
- (iii) At this position it will hit the spring. What is the maximum compression s of the spring?
- (iv) Now the spring is removed. State the equation of motion and give the general solution for small angles ϕ .
- (v) How long does it take for the rod to reach the vertical position for the first time?

3. Exercise

A mass m_1 is performing an circular motion with period T and an radius R . At $t = 0 \text{ s}$ this mass will hit a mass m_2 at rest in a total inelastic collision.

- (i) What would be the angular velocity of the mass m_1 before the collision?
- (ii) Find the velocity of the mass m_1 before the collision.
- (iii) Determine the velocity u of both masses after the collision.
- (iv) How much energy Q is lost in the collision?
- (v) Find the new radius R' if the circular motion is caused by a magnetic field B and the total mass carries a charge q .
- (vi) What would be the new period T' ?

4. Exercise

An electron with charge e and mass m_e enters with velocity v in x -direction an area where an homogeneous electric field E in y -direction is present.

- (i) What voltage has to be applied to a parallel plate capacitor with area $A = \ell^2$, distance d to produce such an electric field? How much charges are needed on the plates?
- (ii) Find the acceleration of the electron?
- (iii) How long does it take to travel a distance s in x -direction?
- (iv) How far and in which direction would it be deflected from its original moving direction?
- (v) What would be then the absolute value of its velocity?
- (vi) Calculate these quantities for $\ell = 1 \text{ km}$, $d = 2 \text{ m}$, $v = 2 \cdot 10^6 \frac{\text{m}}{\text{s}}$, $E = 400 \frac{\text{N}}{\text{C}}$, $e = 1.6 \cdot 10^{-19} \text{ C}$, $m_e = 9.1 \cdot 10^{-31} \text{ kg}$ and $s = 10 \text{ cm}$ ($\epsilon_0 = 8.85 \cdot 10^{-12} \frac{\text{As}}{\text{Vm}}$).

5. Exercise

A rod with mass m is moving with velocity v on two rails towards an inclined plane. The angle of the inclined plane with respect to the horizontal amounts to α and the rails are connected by resistance R at the bottom of the inclined plane. The rod is exposed to gravitational force.

- (i) At which height the velocity will vanish?
- (ii) What distance s on the inclined plane the rod has travelled until it reaches that position?
- (iii) What is the total force acting on the rod on the inclined plane?

Afterwards a magnetic field B is applied perpendicular to the plane.

- (iv) How large will be the terminal current I ?
- (v) Determine the terminal induced voltage U_{ind}
- (vi) Find the final velocity v of the rod.
- (vii) State the power P_F provided by gravity.

6. Exercise

An empty coil with length ℓ , turns N , area A and inner resistance r is connected to two parallel resistors R_1 and R_2 . Initially a magnetic field flux Φ_0 is running through the coil, but is switched off at $t = 0 \text{ s}$.

- (i) What is the maximal current I_0 which is running through the coil?
- (ii) Determine the equivalent resistance of the whole circuit.
- (iii) Derive the power $P_{r,0}$ which is dissipated by the internal resistance of the coil at $t = 0 \text{ s}$?
- (iv) Find the power $P_{p,0}$ which is provided by the coil at $t = 0 \text{ s}$.
- (v) How large is the maximal voltage $U_{R,0}$ that drops at the two resistors?
- (vi) State the maximal currents $I_{1,0}$ and $I_{2,0}$ through the resistors R_1 and R_2 .
- (vii) Give the current $I_2(t)$ through the resistor R_2 with respect to the time.