


1. Exercise:



A stone $m = 1 \text{ kg}$ is thrown from a tower with height $h = 20 \text{ m}$ horizontally with velocity $v_0 = 10 \frac{\text{m}}{\text{s}}$. Gravity is acting along the vertical with $g = 10 \frac{\text{m}}{\text{s}^2}$.

- (a) Take for example the position for $t = 0$. Draw in the diagram all forces acting on the stone.
- (b) What kind of motion performs the body along the x -direction?
 - uniform motion motion with constant acceleration circular motion
- (c) What kind of motion performs the body along the y -direction?
 - uniform motion motion with constant acceleration circular motion
- (d) What are the initial velocities in x and y -direction?

$v_{0,x} =$

$v_{0,y} =$

- (e) How is the distance traveled along the x -direction depending on time?

$$x(t) =$$

- (f) How is the distance traveled along the y -direction depending on time?

$$y(t) =$$

- (g) How is the velocity in x -direction depending on time?

$$v_x(t) =$$

- (h) How is the velocity in y -direction depending on time?

$$v_y(t) =$$

- (i) Solve the equation for $y(t)$ in order to get the time when the stone impacts with the ground.

$$t =$$

- (j) Put this time into the formula for $x(t)$ to get the range:

$$s =$$

- (k) Put the time of impact into the formula of $v_x(t)$ and $v_y(t)$ in order to get the component of the velocity at the time of impact.

$$v_x(t) =$$

$$v_y(t) =$$

- (l) Calculate the absolute value of the velocity at the time of impact.

$$v =$$

- (m) Draw the velocity as well as its components at the time of impact in the diagram above.

- (n) Determine the angle at which the stone impacts with the ground with respect to the horizontal.

$$\alpha =$$